

Table 3-4. Summary of Fish Collected from the Study Area

	Fisherville Pond										Singing Pond		Lake Wildwood	
	MADFW 1992 ^a		Corps 1996 ^b		Corps 1999 ^c		Corps 1999 ^c		Corps 1999 ^c		Corps 1999 ^c		Corps 1999 ^c	
	No. of Fish	Percent of Total	No. of Fish	Percent of Total	North Pool		Central Pool		South Pool		No. of Fish	Percent of Total	No. of Fish	Percent of Total
Black crappie	5	1.4			1	1.1	2	0.74	16	4.8			2	0.87
Bluegill	2	0.6	4	6.3	57	60	197	73	207	62	3	10	169	74
Brown bullhead									1	0.3	13	43.3		
Chain pickerel	2	0.6	2	3.1	1	1.1							2	0.87
Common carp	20	5.6							1	0.3				
Golden shiner	55	15.4	4	6.3			2	0.74	45	13.5				
Largemouth bass	15	4.2	2	3.1	20	21	17	6.3	19	5.7	2	6.7	39	17
Pumpkinseed	2	0.6			10	10.5	44	16.3	41	12.3	5	16.7	4	1.8
Red-fin pickerel											1	3.3		
White perch	3	0.8							2	0.6				
Perch Sp.					1	1.1								
White sucker	210	60	33	51.6							4	13.3		
Yellow bullhead	13	3.6	1	1.6							2	6.7	1	0.44
Yellow perch	27	7.6	18	28.1	5	5.3	8	3	2	0.6			12	5.2
Rainbow trout ^d	1	0.3												
Brook trout ^c	1	0.3												
Total Fish	356		64		95		270		334		30		229	

a. Fish collected by gill net

b. Fish collected by gill net, hoop net, beach seine, and backpack electrofishing equipment

c. Fish collected with boat electrofishing equipment.

d. Fish were stocked holdovers from the Quinsigamond River